O‘ZBEKISTON RESPUBLIKASI

OLIY TA’LIM, FAN VA INNOVATSIYALAR VAZIRLIGI

NAMANGAN DAVLAT UNIVERSITETI

“MARKAZIY OSIYODA BIOLOGIK XILMA-XILLIKNI SAQLASH: MUAMMOLAR, YECHIMLAR VA ISTIQBOLLARI” MAVZUSIDAGI

III XALQARO KONFERENSIYA

Namangan - 2026

ROLE OF Ca^{2+} EXCHANGE IN MUSCLE CONTRACTION AND FATIGUE MECHANISMS IN ATHLETES

S.Z. Omonturdiev[1], A.O. Khasanov[2], L.A. Pulatov[1]
[1]Oriental university, Tashkent, Uzbekistan
[2]Namangan State University, Namangan Region, Uzbekistan
E-mail: siroj.2012@mail.ru

Abstract. Calcium ions (Ca^{2+}) are the primary regulators of excitation–contraction coupling in skeletal muscles. Efficient calcium exchange in athletes ensures high muscle strength, endurance, and rapid recovery. This article analyzes the role of Ca^{2+} ions in muscle contraction, sarcoplasmic reticulum activity, and fatigue mechanisms during physical exercise. Particular attention is given to calcium channels, SERCA pumps, oxidative stress, and training-induced adaptations in skeletal muscles. The findings indicate that regular physical training improves calcium transport efficiency, delays fatigue onset, and enhances athletic performance.

Keywords: sports physiology, calcium ions, skeletal muscle, fatigue, SERCA, sarcoplasmic reticulum, ion channels.

Introduction

Sports physiology investigates the functional and adaptive changes that occur in the human body during physical activity. Modern athletic performance depends not only on organ-level adaptations but also on molecular and cellular mechanisms within skeletal muscles. Among these mechanisms, calcium ions (Ca^{2+}) play a central role in regulating muscle contraction, intracellular signaling, and energy metabolism (Hall & Hall, 2021).

Calcium ions function as universal intracellular messengers that participate in excitation–contraction coupling, enzyme activation, mitochondrial metabolism, and neuromuscular transmission (Hille, 2001). During muscle activation, action potentials propagate along the sarcolemma and T-tubules, triggering Ca^{2+} release from the sarcoplasmic reticulum. Released calcium binds to troponin C and initiates interaction between actin and myosin filaments, resulting in muscle contraction (Berchtold et al., 2000).

Sustained physical exercise requires repeated cycles of calcium release and reuptake. Therefore, efficient Ca^{2+} transport systems are essential for maintaining muscle performance and preventing fatigue. Disturbances in calcium homeostasis during high-intensity exercise are strongly associated with reduced contraction force, metabolic stress, and impaired recovery (Allen et al., 2008).

Recent scientific studies demonstrate that regular physical training induces adaptive changes in calcium handling systems, including increased SERCA pump activity, improved calcium channel sensitivity, and enhanced sarcoplasmic reticulum function (Wilmore et al., 2023). These adaptations contribute to increased endurance and fatigue resistance in athletes.

The aim of this article is to analyze the role of Ca^{2+} exchange in muscle contraction and fatigue mechanisms in athletes from the perspective of sports physiology.

Materials and Methods

This article was prepared based on a review and analysis of scientific literature related to sports physiology, skeletal muscle physiology, calcium signaling, and ion channel function. International review articles, physiological textbooks, and experimental research papers were used as primary scientific sources.

The analysis focused on:

- excitation–contraction coupling mechanisms;
- calcium transport in skeletal muscle fibers;
- sarcoplasmic reticulum function;
- SERCA pumps and ryanodine receptor activity;
- fatigue-related disturbances in calcium homeostasis;
- adaptive changes induced by regular physical training.

Comparative and descriptive physiological approaches were applied to summarize modern scientific understanding regarding calcium regulation in athletes.

Results

Role of Ca^{2+} in excitation–contraction coupling

Muscle contraction begins with propagation of an action potential along the sarcolemma and transverse tubules. Depolarization activates voltage-sensitive dihydropyridine receptors, which subsequently stimulate ryanodine receptors located on the sarcoplasmic reticulum membrane. As a result, Ca^{2+} ions are released into the cytoplasm (Rossi & Dirksen, 2006).

Released calcium binds to troponin C, causing displacement of tropomyosin and exposure of active sites on actin filaments. This process enables actin–myosin cross-bridge formation and force generation (Berchtold et al., 2000).

The relationship between intracellular calcium concentration and contraction force can be represented as:

$$F \propto [Ca^{2+}]_i$$

This relationship demonstrates that muscle force production directly depends on intracellular calcium concentration. Regular physical training improves calcium channel sensitivity and increases the efficiency of excitation–contraction coupling, allowing athletes to produce stronger and faster muscular contractions (Wilmore et al., 2023).

Sarcoplasmic reticulum and SERCA pump activity

The sarcoplasmic reticulum serves as the primary intracellular calcium storage organelle in skeletal muscle fibers. During muscle relaxation, SERCA (Sarcoplasmic/Endoplasmic Reticulum Ca^{2+}-ATPase) pumps actively transport calcium ions from the cytoplasm back into the sarcoplasmic reticulum using ATP energy (Place et al., 2010).

Calcium reuptake can be expressed as:

$$Ca^{2+}_{cytosol} \rightarrow Ca^{2+}_{SR}$$

Efficient SERCA activity is essential for rapid muscle relaxation and repeated contractions during athletic performance. During intense exercise, ATP depletion may impair SERCA pump function, resulting in delayed calcium reuptake and muscle fatigue (Allen et al., 2008).

Training-induced increases in SERCA protein expression have been observed in athletes, suggesting improved recovery capacity and enhanced calcium cycling efficiency (MacIntosh et al., 2012).

Calcium homeostasis and muscle fatigue

High-intensity exercise causes significant metabolic and ionic disturbances within skeletal muscles. ATP depletion, accumulation of lactate and inorganic phosphate, and increased oxidative stress negatively affect calcium homeostasis (Fitts, 2008).

Major mechanisms contributing to muscle fatigue include:

- reduced calcium release from the sarcoplasmic reticulum;
- decreased sensitivity of troponin to Ca^{2+};
- impaired SERCA pump activity;
- oxidative damage to calcium channels;
- disruption of excitation–contraction coupling (Powers & Jackson, 2008).

Reactive oxygen species generated during exhaustive exercise may damage ryanodine receptors and membrane proteins involved in calcium transport. Consequently, calcium regulation becomes impaired, reducing muscle contraction efficiency (Place et al., 2010).

Prolonged elevation of cytoplasmic Ca^{2+} concentration may also activate proteolytic enzymes and contribute to cellular damage in fatigued muscles (Clausen, 2003).

Adaptive changes induced by physical training

Regular exercise training induces important structural and functional adaptations in calcium transport systems. Athletes commonly demonstrate:

- increased sarcoplasmic reticulum volume;
- enhanced SERCA pump activity;
- improved calcium channel sensitivity;
- increased mitochondrial ATP production;
- greater resistance to oxidative stress (Wilmore et al., 2023).




Figure 1. *Ca²⁺* exchange mechanisms in skeletal muscle during excitation–contraction coupling, showing calcium release from the sarcoplasmic reticulum, binding to troponin, actin–myosin cross-bridge formation, and ATP-dependent reuptake via SERCA pumps, along with fatigue-related disruptions and training-induced adaptations in athletes.

Aerobic training particularly improves mitochondrial function and ATP synthesis, supporting more efficient calcium reuptake during repeated contractions (Powers & Jackson, 2008).

Furthermore, long-term training enhances antioxidant defense systems, helping protect ion channels and membrane proteins from oxidative damage. These physiological adaptations contribute to delayed fatigue development and improved athletic performance.

Table 1.

Role of Ca^{2+} Exchange in Skeletal Muscle Function and Fatigue Development

Physiological Component	Main Function	Effect During Exercise	Contribution to Fatigue
Sarcoplasmic Reticulum	Stores and releases Ca^{2+} ions	Increases intracellular Ca^{2+} concentration during contraction	Reduced Ca^{2+} release decreases contraction force
Ryanodine Receptors (RyR)	Mediate Ca^{2+} release from SR	Support excitation–contraction coupling	Oxidative damage impairs Ca^{2+} release
SERCA Pumps	Return Ca^{2+} into SR during relaxation	Accelerate muscle relaxation	ATP depletion reduces pump activity
Troponin C	Binds Ca^{2+} ions	Initiates actin–myosin interaction	Reduced Ca^{2+} sensitivity weakens contraction
Mitochondria	Produce ATP for Ca^{2+} transport	Maintain energy supply during exercise	Reduced ATP synthesis impairs Ca^{2+} homeostasis
Calcium Channels	Regulate Ca^{2+} influx and signaling	Improve muscle activation efficiency	Channel dysfunction contributes to fatigue
Oxidative Stress	Alters ion channel activity	Increases during prolonged exercise	Damages calcium transport proteins

Discussion

The present analysis demonstrates that Ca^{2+} ions are fundamental regulators of skeletal muscle performance in athletes. Efficient calcium release and reuptake ensure rapid contraction–relaxation cycles necessary for sustained physical activity (Hall & Hall, 2021).

Previous investigations have shown that exercise-induced increases in SERCA activity improve muscle relaxation and reduce fatigue development (Allen et al., 2008). Enhanced calcium handling also contributes to improved muscular endurance and recovery capacity.

Disturbances in calcium homeostasis during exhaustive exercise significantly impair muscle performance. ATP depletion and oxidative stress reduce calcium transport efficiency and weaken actin–myosin interactions, resulting in decreased force production (Fitts, 2008).

The findings also suggest that maintaining calcium balance is essential for optimizing athletic performance and preventing exercise-induced muscle dysfunction. Therefore, calcium signaling mechanisms represent an important research area in sports physiology and exercise science.

Conclusion

Calcium ions are essential regulators of contraction and relaxation processes in skeletal muscles. Efficient Ca^{2+} exchange contributes significantly to muscle strength, endurance, and athletic performance. Disturbances in calcium homeostasis are among the major mechanisms responsible for muscle fatigue during intense physical exercise. Regular physical training improves calcium transport systems, enhances SERCA pump activity, and increases resistance to fatigue. Understanding calcium signaling mechanisms is important for sports physiology and may contribute to the development of more effective training and recovery strategies for athletes.

References

1. Allen, D. G., Lamb, G. D., & Westerblad, H. (2008). Skeletal muscle fatigue: Cellular mechanisms. *Physiological Reviews, 88*(1), 287–332.
2. Berchtold, M. W., Brinkmeier, H., & Müntener, M. (2000). Calcium ion in skeletal muscle: Its crucial role for muscle function. *Physiological Reviews, 80*(3), 1215–1265.
3. Clausen, T. (2003). Excitation-induced exchange of Na^+, K^+, and Ca^{2+} in skeletal muscle. *Physiological Reviews, 83*(4), 1269–1324.
4. Fitts, R. H. (2008). The cross-bridge cycle and skeletal muscle fatigue. *Journal of Applied Physiology, 104*(2), 551–558.
5. Hall, J. E., & Hall, M. E. (2021). *Guyton and Hall textbook of medical physiology* (14th ed.). Elsevier.
6. Hille, B. (2001). *Ion channels of excitable membranes* (3rd ed.). Sinauer Associates.
7. MacIntosh, B. R., Holash, R. J., & Renaud, J. M. (2012). Skeletal muscle fatigue: Regulation of excitation–contraction coupling to avoid metabolic catastrophe. *Journal of Cell Science, 125*(9), 2105–2114.
8. Place, N., Yamada, T., Bruton, J. D., & Westerblad, H. (2010). Interdependence of calcium handling and metabolism in skeletal muscle during fatigue. *Journal of Physiology, 588*(15), 2865–2875.
9. Powers, S. K., & Jackson, M. J. (2008). Exercise-induced oxidative stress: Cellular mechanisms and impact on muscle force production. *Physiological Reviews, 88*(4), 1243–1276.
10. Rossi, A. E., & Dirksen, R. T. (2006). Sarcoplasmic reticulum: The dynamic calcium governor of muscle. *Muscle & Nerve, 33*(6), 715–731.
11. Wilmore, J. H., Costill, D. L., & Kenney, W. L. (2023). *Physiology of sport and exercise* (8th ed.). Human Kinetics.

PHLOMOIDES STELLATA VA *PHLOMOIDES KURPSAICA* (LAMIACEAE) O‘SIMLIKLARINING GERBARIY TAHLILI

Sh.G‘. Mavlonova
Namangan davlat universiteti magistranti
E-mail: shaxnozamavlonova64@gmail.com

Abstract. *Phlomoides stellata* is a perennial plant species belonging to the Lamiaceae family and distributed in the mountainous regions of Central Asia. This species is characterized by its thick rhizome, leaves covered with stellate hairs, and whitish flowers. The stems are erect, while the leaves are cordate at the base with rounded dentate margins. The inflorescence consists of 3-5 verticillasters . The species mainly occurs on rocky and calcareous slopes of the Turkestan Range in Tajikistan. According to its morphological characteristics, it is closely related to *Phlomoides cordifolia*, but differs in the indumentum of the leaves, the structure of the inflorescence, and several features of the generative organs. This species is of considerable scientific importance in botanical, floristic, and herbarium studies.

Keywords: *Phlomoides stellata, Phlomoides cordifolia, Phlomoides kurpsaica*, Lamiaceae, perennial plant, herbarium analysis, Central Asia flora, Turkestan Range, morphology, stellate hairs, botanical taxonomy.

Kirish

218	**TENEBRIONIDAE OILASINING YEVROPA MAMLAKATLARIDA O‘RGANILISH TARIXI VA ZAMONAVIY TADQIQOTLAR TAHLILI** A.A. Ma’rupov, S.A. Asqaraliyeva……	706
219	**ATROF-MUHITNI KO‘KALAMZORLASHTIRISHDA PAEONIA SUFFRUTICOSA NAVLARINING O‘RNI** F.F. Nurmuxamedova, X.Q. Haydarov……	708
220	**BUXORO VILOYATI SHAROITIDA AQLLI SHAHAR KONSEPSIYASINI JORIY ETISHDA EKOLOGIK TEXNOLOGIYALARNING EKOLOGIK-IQTISODIY SAMARADORLIGI** Sulaymonova Mastura O‘tkir qizi, Qobulova Barno Baxriddin qizi……	710
221	**ROLE OF Ca^{2+} EXCHANGE IN MUSCLE CONTRACTION AND FATIGUE MECHANISMS IN ATHLETES** S.Z. Omonturdiev, A.O. Khasanov, L.A. Pulatov……	713
222	**PHLOMOIDES STELLATA VA PHLOMOIDES KURPSAICA (LAMIACEAE) O‘SIMLIKLARINING GERBARIY TAHLILI** Sh.G‘. Mavlonova……	716
223	**NOYOB DORIVOR O‘SIMLIK DIANTHUS UZBEKISTANICUS URUG‘INING MORFOLOGIYASI VA UNISH BIOLOGIYASI** Turopova Muxlisa Ilhom qizi……	719
224	**YOVVOYI CHO’L MUSHUGI - FELIS LIBICANING TOG’ MINTAQALARI BO’YLAB VERTIKAL MIGRATSIYASI** Barakayev S.M., Tovboyeva O.O……	722
225	**QANDLI DIABET BILAN OG'RIGAN AYOLLARNING HOMILADORLIK DAVRIDA AZOT OKSIDINING ROLI** Kabirova Zarnigor Akmaljon qizi……	725
226	**ASALARI PARAZITLARINING TARQALISHI VA ASALARICHILIK MAHSULDORLIGIGA TA’SIRI** M.F. Mirzaxmedov……	729
227	**FARG‘ONA VODIYSI BIOXILMA-XILLIGI: STRESS OMILLAR, ANTROPOGEN TA’SIR VA SAQLASH STRATEGIYALARI** D.J.Komilov, R.A Rahimjonova……	733
228	**ZIG‘IR GERMOPLAZMASIDA YUQORI MOYDORLIK KO‘RSATKICHLARI BILAN ASSOTSIATSIYALANGAN DNK MARKERLARNING MOLEKULYAR-GENETIK TAHLILI** U.Sh. Bakhadirov, N.Kh.Khudoyberdiev, U. Yuldashev, A.A. Dolimov……	736
229	**BUG‘DOY NAVLARINING SARIQ ZANG KASALLIGIGA CHIDAMLILIGINI FENOTIPIK BAHOLASH** Shokirova D.Sh., To‘raqulov X.S., Meliev S.K., Aytenov I.S., Ochilov B.O., Xalilellayeva G.O., Murodova S.M., Mardonova M.R.……	740
230	**ЗАГРЯЗНЕНИЕ СТОЧНЫХ ВОД В ЧАТКАЛЬСКОМ РЕГИОНЕ И ЕГО ВЛИЯНИЕ НА ЗДОРОВЬЕ** Мырзабаев Иманкул Тургунбекович……	743